arXiv:math/9905001v1 [math.AG] 1 May 1999

Tacnodes and cusps

Joaquim Roé *
Departament d'Àlgebra i Geometria, Universitat de Barcelona,
Gran Via, 585, E-08007, Barcelona.
e-mail: jroevell@cerber.mat.ub.es

February 1, 2008

1 Introduction

The study of linear systems of algebraic plane curves with fixed imposed singularities is a classical subject which has recently experienced important progress. The Horace method introduced by A. Hirschowitz in [23] has been successfully exploited to prove many H^1–vanishing theorems, even in higher dimension. Other specialization techniques, which include degenerations of the plane, are due to Z. Ran [32], [33] and C. Ciliberto and R. Miranda [8]. In [14], G. M. Greuel, C. Lossen and E. Shustin use a local specialization procedure together with the Horace method to give the first asymptotically proper general existence criterion for singular curves of low degree. In this paper we develop a specialization method which allows us to compute the dimension of several linear systems as well as to substantially improve the bounds of [14] for curves with tacnodes and cusps.

A closed subscheme $Z \subset \mathbb{P}^2$ is said to have *maximal rank in degree* d if the canonical map $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(d)) \to H^0(Z, \mathcal{O}_Z(d))$ has maximal rank (cf. [23]). If Z is a zero–dimensional scheme of length N, this means that either there are no curves of degree d containing Z or Z imposes N independent linear conditions to curves of degree d. If Z has maximal rank in all degrees then we say simply that it has maximal rank.

For some classes of zero-dimensional schemes, it is known that general members have maximal rank. For example, a general union of double points has maximal rank. In many other cases, however, a maximal rank statement has been conjectured only (cf. [18], [24], [28]). We will consider schemes $Z = Z_1 \cup Z_2 \cup \ldots \cup Z_\rho$ where Z_i are unibranched cluster schemes all whose points but one, whether proper or infinitely near, are taken with multiplicity ≤ 2. In theorem 4.8 we prove that, under some mild numerical conditions, a scheme Z as above whose points are in general position has maximal rank. This generalizes and unifies a range of previously known results:

- In [23], A. Hirschowitz proves that a general union of distinct points with multiplicity 2 or 3 has maximal rank. Our result generalizes the multi-

*Partially supported by CIRIT 1997FI-00141, CAICYT PB95-0274, and "AGE-Algebraic Geometry in Europe" contract no. ERB940557.

plicity 2 case by allowing infinitely near points, which are not easily dealt with by the Horace method.

- M. V. Catalisano and A. Gimigliano in [6] and C. Ciliberto and R. Miranda in [7] prove that a union of general unibranched cluster schemes all whose points have multiplicity one (curvilinear schemes) has maximal rank. Our result generalizes this by allowing multiplicity 2 points.

- In [8], C. Ciliberto and R. Miranda consider quasi-homogeneous schemes, which consist of distinct multiple points, all whose multiplicities but one are taken to be equal. They prove a quite general maximal rank theorem which includes the computation of the superabundant systems (corresponding to multiplicities for which the general scheme does not have maximal rank). In the particular case in which one has a point of multiplicity not bigger than 5 and all other points have multiplicity 2, we extend their result by allowing infinitely near points.

Other results concerning linear systems with infinitely near base points can be found in the literature, mainly in [17], [20], [14] and [27]. Harbourne's results deal with clusters of points lying on conics and cubics, which is not necessary to us, and the H^1–vanishing of [14] and [27] is weaker than proving maximal rank.

The maximal rank theorem will allow us to prove the existence of irreducible curves of low degree with tacnodes and higher order cusps. The reasoning is similar to that of [2], [14] or [27]. Theorem 4.12 is much sharper than the result by Greuel, Lossen and Shustin (which nevertheless applies to any kind of plane singularity) and also sharper than the one Lossen obtains for tacnodes and cusps (A-singularities), in part because our specialization avoids the use of the Viro method to "glue" the singularities. On the other hand, Barkats' result is a little bit sharper than ours, but it is restricted to nodes and ordinary cusps only.

2 Preliminaries

Let k be an algebraically closed field of characteristic zero, p a smooth point of a surface S defined over k, $\mathcal{O} = \mathcal{O}_{S,p}$ the local ring of p on S, $\mathfrak{m} = \mathfrak{m}_p$ its maximal ideal.

Consider a sequence of blowing-ups

$$S_r \xrightarrow{\sigma_r} S_{r-1} \longrightarrow \cdots \xrightarrow{\sigma_2} S_1 \xrightarrow{\sigma_1} S_0 = S$$

where σ_1 is the blowing-up of p and for $i > 1$ the center of σ_i is a point p_i which lies on the exceptional divisor of σ_{i-1}. The sequence $K = (p_1 = p, p_2, \ldots, p_r)$ is a cluster with origin at p for which every point is infinitely near the preceding one; we will call these clusters *unibranched*. We write $S_K = S_r$ and $\pi_K : S_K \to S$ the composition of the blowing-ups. Usual facts known for clusters hold in particular for unibranched clusters; we now review some of them, referring the reader to [4], [5] for the proofs.

A point p_j is said to be *proximate* to p_i, $j > i$ if and only if it lies on the exceptional divisor of blowing up p_i (that is, $j = i + 1$) or on its *strict* transform when $j > i + 1$. Every point in a unibranched cluster is proximate either to one or to two points, except for p_1, which is proximate to no one; if p_j is proximate to p_i then all the points between them are also proximate to p_i.

If p_j is proximate to p_i then there is a unique point in E_j proximate to both p_i and p_j; otherwise there is none. If a point is proximate to two points, it is called a *satellite*, otherwise it is *free*. When considering more than one cluster at a time, we will write $p_i(K)$ and $S_i(K)$ for the i–th point of the cluster K and the surface obtained by blowing up the first i points of K.

A *system of multiplicities* for a cluster $K = (p_1, p_2, \dots, p_r)$ is a sequence of integers $\mathbf{m} = (m_1, m_2, \dots, m_r)$, and a pair $(K, \mathbf{m})$ where K is a cluster and $\mathbf{m}$ a system of multiplicities is called a *weighted cluster*. A system of multiplicities like

$$(m, 2, 2, \overset{i}{\breve{\dots}}, 2, 1, 1, \overset{j}{\breve{\dots}}, 1, 0, 0, \dots, 0)$$

will be designed as $(m, 2^i, 1^j)$. Given a weighted cluster, there are an ideal and a zero–dimensional subscheme of S associated to it. Let E_i be the pullback (total transform) in S_K of the exceptional divisor of blowing up p_i. Then the ideal sheaf

$$\mathcal{H}_{K,\mathbf{m}} = (\pi_K)_* \, \mathcal{O}_{S_K}(-m_1 E_1 - m_2 E_2 - \dots - m_r E_r)$$

is supported at p, its stalk at p is a complete $\mathfrak{m}$–primary ideal $H_{K,\mathbf{m}} \subset \mathcal{O}$ and defines a zero–dimensional subscheme $Z_{K,\mathbf{m}}$ of S. For K unibranched we call $Z_{K,\mathbf{m}}$ a *unibranched cluster scheme*. As an aside, note that if $I \subset \mathcal{O}$ is a complete $\mathfrak{m}$–primary ideal then there is a weighted cluster $(K, \mathbf{m})$ such that $I = H_{K,\mathbf{m}}$, but this cluster does not need to be unibranched.

The same (unibranched) cluster scheme is sometimes defined by different (unibranched) weighted clusters. In this case we will say that both clusters are equivalent. In order to have a well defined weighted cluster associated to every cluster scheme, one considers the notion of *consistent* clusters, which we define next. The *proximity inequality* at p_i is

$$m_i \geq \sum_{p_j \text{ prox. to } p_i} m_j.$$

A weighted cluster $(K, \mathbf{m})$ is consistent if and only if it satisfies the proximity inequalities at all its points. Given a (unibranched) cluster scheme Z, there is a unique consistent weighted cluster $(K, \mathbf{m})$ such that $Z = Z_{K,\mathbf{m}}$ and $m_i > 0$ for all i. Furthermore, for a weighted cluster $(K, \mathbf{m})$ non necessarily consistent,

$$\operatorname{length} Z_{K,\mathbf{m}} = \dim \frac{\mathcal{O}}{H_{K,\mathbf{m}}} \leq \sum_{i=1}^{r} \frac{m_i \, (m_i + 1)}{2},$$

with the equality holding if $(K, \mathbf{m})$ is consistent.

Given an arbitrary weighted cluster $(K, \mathbf{m})$ there is a procedure called *unloading* (see [5, 4], [10, IV.II], or [4]) which gives a new system of multiplicities $\delta(\mathbf{m}) = \delta_K(\mathbf{m})$ such that $(K, \delta(\mathbf{m}))$ is consistent and equivalent to $(K, \mathbf{m})$. In each step of the procedure, one *unloads* some amount of multiplicity on a point p_i whose proximity inequality is not satisfied, from the ponts proximate to it. This means that there is an integer $n > 0$ such that, increasing the multiplicity of p_i by n and decreasing the multiplicity of every point proximate to p_i by n, the resulting weighted cluster is equivalent to $(K, \mathbf{m})$ and satisfies the proximity inequality at p_i. In other words, if $\tilde{E}_i \subset S_K$ is the strict transform of the exceptional divisor of blowing-up p_i, $D = -m_1 E_1 - m_2 E_2 - \dots - m_r E_r$ and and

$\tilde{E}_i \cdot D < 0$ then one chooses n as the minimal integer with $\tilde{E}_i \cdot (D - n\tilde{E}_i) \geq 0$ and replaces D by $D - n\tilde{E}_i$. A finite number of unloading steps lead to the desired equivalent consistent cluster $(K, \delta(\mathbf{m}))$.

Let $C \subset S$ be a curve (more generally, a divisor). For any proper or infinitely near point q of S we write $e_q(C)$ the multiplicity of (the strict transform of) C at q. Let $(K, \mathbf{m})$ be a weighted cluster of r points and $i \leq r$. Let S_i be the surface obtained by blowing up the first i points of K, $\bar{C} \subset S_i$ the pullback of C and for $1 \leq j \leq i$, $E_j \subset S_i$ the pullback of the exceptional divisor of blowing up $p_j(K)$. We define the *virtual transform* of C in S_i relative to the system $\mathbf{m}$ as the divisor

$$\tilde{C} = \bar{C} - m_1 E_1 - m_2 E_2 - \cdots - m_i E_i \ .$$

A curve C contains the cluster scheme $Z_{K,\mathbf{m}}$ if and only if its virtual transform in S_{i-1} has multiplicity at least m_i at p_i for all i. Then we say that C *goes through* $(K, \mathbf{m})$. If $e_{p_i} = m_i$ for all i then the virtual transform coincides with the strict transform; if furthermore $\bar{C}$ is a divisor with normal crossings then we say that C goes *sharply* through $(K, \mathbf{m})$. In this case, π_K is an embedded resolution of singularities for C, and $(K, \mathbf{m})$ determines the equisingularity class of C. There are curves going sharply through $(K, \mathbf{m})$ if and only if $(K, \mathbf{m})$ is consistent, and in this case they are all equisingular.

We define next varieties $Y_i \subset X_i$ and smooth surjective morphisms $\psi_i : X_i \to Y_{i-1}$ of relative dimension 2, as follows: Let $Y_{-1} = \operatorname{Spec} k$, $X_0 = S$ and $Y_0 = \{p\}$, and for $i > 0$, let

$$X_i \xrightarrow{b_i} Y_{i-1} \times_{Y_{i-2}} X_{i-1}$$

be the blowing-up along the diagonal $\Delta(Y_{i-1}) \subset Y_{i-1} \times_{Y_{i-2}} X_{i-1}$, let $Y_i \subset X_i$ be the exceptional divisor, and ψ_i the composition $\pi_{Y_{i-1}} \circ b_i : X_i \to Y_{i-1}$. We define also morphisms $\pi_i = \pi_{X_{i-1}} \circ b_i$ and $\pi_{i,j} = \pi_{j+1} \circ \cdots \circ \pi_{i-1} \circ \pi_i : X_i \to X_j$.

Remark that by construction the variety Y_i is irreducible and smooth, and $\psi_i|_{Y_i} : Y_i \to Y_{i-1}$ is a $\mathbb{P}^1$–bundle, therefore Y_i is also projective and rational for all i. The varieties X_i are irreducible, smooth, projective and rational if S is. Y_{r-1} can be identified with the set of all unibranched clusters of r points of S with origin at p (cf. [34], [19], [31]) in such a way that for all $K = (p_1, p_2, \ldots, p_r) \in Y_{r-1}$, denoting by $j_K : \{K\} \to Y_{r-1}$ the inclusion one has a pullback diagram

$$\begin{array}{ccccc} S_K & & \xrightarrow{\quad i_K \quad} & & X_r \\ \downarrow & & & & \downarrow{\scriptstyle \psi_r} \\ \operatorname{Spec} k & = & \{K\} & \xrightarrow{j_K} & Y_{r-1} \end{array}$$

where i_K is a closed immersion, and a commutative square

$$\begin{array}{ccc} S_K & \xrightarrow{\quad i_K \quad} & X_r \\ {\scriptstyle \sigma_r}\downarrow & & \downarrow{\scriptstyle \pi_r} \\ S_{\breve{K}} & \xrightarrow{i_{\breve{K}}} & X_{r-1} \end{array}$$

where $\breve{K} = (p_1, p_2, \ldots, p_{r-1}) = \psi_{r-1}(K) \in Y_{r-2}$ is the cluster of the first $r-1$ points of K, and σ_r is the blowing-up of $p_r(K)$. Remark that $S_{\breve{K}} = S_{r-1}(K)$ and $i_{\breve{K}}(p_r(K)) = K$.



Figure 1: Enriques diagram of a cluster in U_s

For every pair of integers s, t such that $r \geq s > t \geq 1$ the subset of Y_{r-1} containing exactly the clusters K with $p_s(K)$ proximate to $p_t(K)$ is an irreducible closed subvariety $P_{s,t}$ (cf. [34]). There are also open and dense subsets $U_{s,t} \subset P_{s,t}$ which contain all unibranched clusters with $p_i(K)$ proximate to $p_{i-1}(K)$ for all $i > 1$ and to $p_t(K)$ if $s \geq i > t$, and no other proximity relations. We will write $P_s = P_{s,1}$ and $U_s = U_{s,1}$. We say that a system of multiplicities $\mathbf{m}$ is consistent in a subset $W \subset Y_r$ when the weighted cluster $(K, \mathbf{m})$ is consistent for all $K \in W$.

Lemma 2.1. *Fix a cluster $K = (p_1, p_2, \dots, p_r) \in U_s$. Let i, j be positive integers such that $r = i + j + 1$ and consider the systems of multiplicities*

$$\mathbf{m}_- = (m_1, 2^i, 1^j)$$
$$\mathbf{m}_+ = (m_1, 2^{i+1}, 1^{j-1}) .$$

Let $I \subset \mathcal{O}$ be an ideal such that

$$H_{K,\mathbf{m}_+} \subsetneq I \subsetneq H_{K,\mathbf{m}_-} .$$

Then there is a point $q \in E_r$ such that $I = H_{K_q,\mathbf{m}_0}$ with $K_q = (p_1, p_2, \dots, p_r, q)$ and $\mathbf{m}_0 = (m_1, 2^i, 1^{j+1})$.

Proof. For any $q \in E_r$ let q' be the unique point that is proximate to q and to p_r, and consider $K'_q = (p_1, p_2, \dots, p_r, q, q')$ and $\mathbf{m}' = (m_1, 2^i, 1^{j+2})$. By performing successive unloading steps on $p_r, p_{r-1}, \dots, p_{i+2}$ we see that $(K'_q, \mathbf{m}')$ is equivalent to $(K, \mathbf{m}_+)$ and since $(K'_q, \mathbf{m}')$ results from $(K, \mathbf{m}_-)$ by adding two simple points,

$$\dim \frac{H_{K,\mathbf{m}_-}}{H_{K,\mathbf{m}_+}} \leq 2 .$$

The hypothesis on I implies that the inequality is in fact an equality and

$$\dim \frac{I}{H_{K,\mathbf{m}_+}} = \dim \frac{H_{K,\mathbf{m}_-}}{I} = 1 ,$$

so $I = H_{K,\mathbf{m}_+} + (f)$ for some $f \in H_{K,\mathbf{m}_-} \setminus H_{K,\mathbf{m}_+}$. f is the equation of a germ of curve ξ which goes through $(K, \mathbf{m}_-)$ but not through $(K, \mathbf{m}_+)$, therefore its virtual transform at p_r relative to the system of multiplicities $\mathbf{m}_-$ is smooth, so it has a unique point $q \in E_r$. It is clear that $I = H_{K,\mathbf{m}_+} + (f) \subseteq H_{K_q,\mathbf{m}_0}$ and $\dim\left(H_{K_q,\mathbf{m}_0}/H_{K,\mathbf{m}_+}\right) \leq 1$, so $I = H_{K_q,\mathbf{m}_0}$. □

Proposition 2.2. *Let $(K, \mathbf{m})$ be a weighted cluster supported at p and $f \in \mathcal{O}$ the equation of a germ of curve ξ through p. Write $e_i = e_{p_i}(\xi)$. Then the conductor $(H_{K,\mathbf{m}} : f)$ is $H_{K,\mathbf{m}'}$ with*

$$m_i' = m_i - e_i\,.$$

Proof. The total transform of ξ in S_K is

$$\tilde{\xi} + e_1E_1 + e_2E_2 + \cdots + e_rE_r$$

where $\tilde{\xi}$ is the strict transform. Let $g \in \mathcal{O}$ be the equation of a second germ η. If $g \in H_{K,\mathbf{m}'}$ the total transform of $\xi + \eta$ is

$$\tilde{\xi} + C + ((m_1 - e_1) + e_1)E_1 + ((m_2 - e_2) + e_2)E_2 + \cdots + ((m_r - e_r) + e_r)E_r$$

with C effective, so clearly $fg \in H_{K,\mathbf{m}}$. Conversely, if $fg \in H_{K,\mathbf{m}}$, then the total transform of $\xi + \eta$ is

$$\tilde{\xi} + C + m_1E_1 + m_2E_2 + \cdots + m_rE_r$$

with C effective, so the total transform of η must be

$$C + (m_1 - e_1)E_1 + (m_2 - e_2)E_2 + \cdots + (m_r - e_r)E_r$$

and $g \in H_{K,\mathbf{m}'}$. □

If $Z \subset S$ is a zero–dimensional scheme defined by the ideal sheaf $\mathcal{I}_{Z/S}$ and $C \subset S$ is a curve, then there is an exact sequence

$$0 \longrightarrow \mathcal{I}_{Z'/S}(-C) \longrightarrow \mathcal{I}_{Z/S} \longrightarrow \mathcal{I}_{(Z\cap C)/C} \longrightarrow 0$$

where $\mathcal{I}_{(Z\cap C)/C} = \mathcal{I}_{Z/S} \otimes \mathcal{O}_C$ and $\mathcal{I}_{Z'/S}$ defines a zero–dimensional scheme $Z' \subset S$. In the context of the Horace method, it is usually called the *residual exact sequence* and Z' is the *residual scheme* of Z with respect to C (cf. [23]).

Corollary 2.3. *If $Z_{K,\mathbf{m}} \subset S$ is a cluster scheme and C is a curve with $e_{p_i}(C) = e_i$ then the residual scheme Z' of Z with respect to C is the cluster scheme $Z_{K,\mathbf{m}'}$ with $\mathbf{m}'$ as in proposition 2.2.*

Proposition 2.4. *Given a cluster $K = (p_1, p_2, \ldots, p_r) \in U_s$ and a point $q \in E_r$, consider $K_q = (p_1, p_2, \ldots, p_r, q)$. Suppose that $r \geq (s-1)(s-2)$. Then for any $q, q' \in E_r$, each of them proximate only to p_r, there is an automorphism $\varphi^* : \mathcal{O} \to \mathcal{O}$ such that, for any system of multiplicities $\mathbf{m}$,*

$$\varphi^*(H_{K_q,\mathbf{m}}) = H_{K_{q'},\mathbf{m}}\,.$$

In particular, $Z_{K_q,\mathbf{m}} \cong Z_{K_{q'},\mathbf{m}}$.

Proof. Let $\mathbf{m}_0 = (s-1, 1^r)$. It will be enough to see that there are open neighbourhoods $V_i \subset S_{i-1}$ of p_i and $V_{r+1} \subset S_K$ containing both q and q', and isomorphisms $\varphi_i : V_i \to V_i$ commuting with the blowing-ups, such that $\varphi_i(p_i) = p_i$ and $\varphi_{r+1}(q) = q'$. This is equivalent to prove that there are isomorphic unibranched germs of curve ξ and ξ' going sharphly through $(K_q, \mathbf{m}_0)$ and $(K_{q'}, \mathbf{m}_0)$ respectively, because then there is a neighbourhood V of p_1 where both ξ and ξ' have a representative and an automorphism φ of V sending one to the other and which therefore lifts to the desired φ_i. Let $\xi : f = 0$ and $\zeta : g = 0$ be arbitrary unibranched germs of curve going sharphly through $(K_q, \mathbf{m}_0)$ and $(K_{q'}, \mathbf{m}_0)$. They are equisingular with a single characteristic exponent $s/(s-1)$ and their intersection multiplicity at p is $\nu \geq (s-1)^2 + (s-1)(s-2) - 1$, so there is an automorphism of the completion

$$\psi^* : \hat{\mathcal{O}} \to \hat{\mathcal{O}}$$

with $\psi^*(f) = g$ (cf. [37]). Let $x, y \in \mathcal{O}$ be a system of parameters; we have $\hat{\mathcal{O}} \cong k[[x, y]]$, and ψ^* can be described by

$$\psi^*(x) = \sum_{i,j>0} a_{ij} x^i y^j$$
$$\psi^*(y) = \sum_{i,j>0} b_{ij} x^i y^j$$

Choose $e \in \mathbb{Z}_{>1}$ such that $\mathfrak{m}^e \subset H_{K_{q'}, \mathbf{m}_0}$. Then

$$\varphi^*(x) = \sum_{i+j \leq e} a_{ij} x^i y^j$$
$$\varphi^*(y) = \sum_{i+j \leq e} b_{ij} x^i y^j$$

define an automorphism of $\mathcal{O}$ such that $\varphi^*(f) = 0$ is the equation of an irreducible germ ξ' going sharply through $(K_{q'}, \mathbf{m}_0)$. Indeed, $f \in \mathfrak{m} = (x, y)$ so $\varphi^*(f) - \psi^*(f) \in \mathfrak{m}^e \subset H_{H_{q'}, \mathbf{m}_0}$ and we know also that $\psi^*(f) \in H_{H_{q'}, \mathbf{m}_0}$. Therefore $\varphi^*(f) = 0$ is the equation of a germ ξ' going through $(K_{q'}, \mathbf{m}_0)$. As φ^* is an automorphism, ξ' is irreducible and goes sharply through $(K_{q'}, \mathbf{m}_0)$. □

3 Specializing unibranched cluster schemes

The subscheme of the Hilbert scheme which parametrizes cluster schemes of a given type, and its relation to the corresponding variety of clusters, has been studied by many authors ([29], [30], [31], [15], among others). However, they usually assume that the proximity relations between points of the clusters remain constant. Little seems to be known about the relative position of these subschemes. Evain computed in [11] and [12] several collisions of points, including all cases with 3 points; this is equivalent to the determination of the closure of the corresponding subschemes of the Hilbert scheme, that is, the specializations of cluster schemes when new proximity relations arise. In this section we show some flat families of unibranched cluster schemes in which proximities vary; we introduce them in order to prove theorems 4.8 and 4.12 but they can have also some interest on their own.

Let x, y be local parameters for $\mathcal{O}$. The inclusion $\mathcal{O} \subset \hat{\mathcal{O}} \cong k[[x, y]]$ allows to write any $f \in \mathcal{O}$ as a formal power series

$$f = \sum_{i,j \geq 0} a_{ij} x^i y^j$$

in a unique way. Thus any polynomial $\Pi(X_{ij}) \in R = k[X_{ij}]_{i,j \geq 0}$ determines a function $\mathcal{O} \to k$, by evaluating at a_{ij}: $\Pi(f) = \Pi(a_{ij})_{i,j \geq 0}$, and to every $f \in \mathcal{O}$ corresponds a maximal ideal

$$\mathfrak{m}_f = \{\Pi \in R \mid \Pi(f) = 0\} \subset R$$

with quotient field equal to k. So we have a mapping $\mathcal{O} \to \operatorname{MaxSpec} R \subset \operatorname{Spec} R$, and it is easy to see that it is injective. For any variety Y we have therefore $\mathcal{O} \times Y \hookrightarrow \operatorname{Spec} R \times Y$, and we will take the Zariski topology on $\mathcal{O} \times Y$ induced by the one on $\operatorname{Spec} R \times Y$. It is easy to see that this Zariski topology does not depend on the local parameters chosen for $\mathcal{O}$. We will prove in 3.2 below that for any system of multiplicities $\mathbf{m}$ the set

$$H_{\mathbf{m}} := \{(f, K) \in \mathcal{O} \times Y_{r-1} \mid f \in H_{K,\mathbf{m}}\}.$$

is Zariski–closed in $\mathcal{O} \times Y_{r-1}$.

Fix for the rest of the section a system of multiplicities $\mathbf{m} = (m_1, m_2, \ldots, m_r)$ which we will apply to any cluster of r points; write $\breve{\mathbf{m}} = (m_1, m_2, \ldots, m_{r-1})$. For a cluster $K = (p_1, p_2, \ldots, p_r)$, we write also $\breve{K} = (p_1, p_2, \ldots, p_{r-1})$.

Lemma 3.1. *For every $K_0 \in Y_{r-1} \subset X_{r-1}$ there are an open neighbourhood $V \subset X_{r-1}$ of K, an isomorphism of k–varieties*

$$U := V \cap Y_{r-1} \xrightarrow{\lambda} k^{r-1} ,$$

two functions

$$x, y \in \Gamma\left(U \times_{Y_{r-2}} V,\ \mathcal{O}_{Y_{r-1} \times_{Y_{r-2}} X_{r-1}}\right)$$

generating the ideal of $\Delta(U)$ and polynomials $A^r_{ij} \in R[u_1, u_2, \ldots, u_{r-1}]$ for $i, j \geq 0$, such that for every $K \in U$ and $f \in H_{K,\mathbf{m}}$ the formal power series

$$\sum_{i,j \geq 0} A^r_{ij}(f, \lambda(K))(x \circ i_{\breve{K}})^i (y \circ i_{\breve{K}})^j \in \mathcal{O}_{S_{r-1}, p_r}$$

and is a local equation for the virtual transform of $f = 0$ at the last point p_r of K.

Remark that the conditions on x, y and U imply that for any cluster $K \in U$ the functions $x, y, \lambda_1 \circ \pi_U - \lambda_1(K), \lambda_2 \circ \pi_U - \lambda_2(K), \ldots, \lambda_{r-1} \circ \pi_U - \lambda_{r-1}(K)$ are a system of parameters for the local ring of (K, K) in $U \times_{Y_{r-2}} V$, the surface

$$\{K\} \times_{Y_{r-2}} X_{r-1} = \{K\} \times i_{\breve{K}}(S_{\breve{K}}) \cong S_{r-1}(K)$$

is locally defined by the equations $\lambda_i \circ \pi_U = \lambda_i(K)$, and $x \circ i_{\breve{K}}$, $y \circ i_{\breve{K}}$ are local parameters for $\mathcal{O}_{S_{r-1}, p_r}$ (recall that $S_{\breve{K}} = S_{r-1}(K)$ and $i_{\breve{K}}(p_r) = K$).

Proof. We proceed by induction on r. For $r = 1$, choose a system of parameters $x, y \in \mathcal{O}$ and a neighbourhood V of p where x and y have regular representatives (which abusing notation we call x, y also) and $V(x, y) \cap V = \{p\}$. Then the claim is clear, because $Y_0 = \{p\}$, and the virtual transform of f at p is f itself, so we can take $A^1_{ij} = X_{ij}$. Suppose now $r > 1$ and apply the induction hypothesis to $(\breve{K}, \breve{\mathbf{m}})$. We obtain the existence of $\breve{V} \subset X_{r-2}$, $\breve{U} = \breve{V} \cap Y_{r-2} \cong k^{r-2}$, $\breve{\lambda}$, $\breve{x}, \breve{y}$, and the polynomials A^{r-1}_{ij} as in the claim. By making a linear substitution in $\breve{x}, \breve{y}$ we may assume that the last point p_r of K_0 lies in the direction of $\breve{y} \circ i_{\breve{K}_0} = 0$. $b^{-1}_{r-1}(\breve{U} \times_{Y_{r-3}} \breve{V})$ is open in X_{r-1}, and is the blowing-up of $\Delta(\breve{U})$ in $\breve{U} \times_{Y_{r-3}} \breve{V}$, which can be described as the subvariety of

$$\breve{U} \times_{Y_{r-3}} \breve{V} \times \mathbb{P}^1$$

given by the equation $u\breve{x} - v\breve{y} = 0$, where $(u : v)$ are projective coordinates of $\mathbb{P}^1$. The exceptional divisor $Y_{r-1} \cap b^{-1}_{r-1}(\breve{U} \times_{Y_{r-3}} \breve{V})$ has equations $\breve{x} = \breve{y} = 0$. We define now V to be the open subset determined by $v \neq 0$, and the isomorphism

$$\lambda : U \longrightarrow k^{r-1} = k^{r-2} \times k$$

as $\lambda = (\breve{\lambda} \circ \psi_{r-1}) \times (u/v)$. Let π_V, π_U be the projections of $V \times_{\breve{U}} U$ on its two factors. Then it is easy to see that

$$\begin{aligned} x &= \breve{x} \circ \pi_V \\ y &= \frac{u}{v} \circ \pi_V - \frac{u}{v} \circ \pi_U = \frac{\breve{y}}{\breve{x}} \circ \pi_V - \lambda_r \circ \pi_U \end{aligned}$$

generate the ideal of $\Delta(U)$.

Because of the induction hypothesis we know that for any $K \in U$ and $f \in H_{K,\mathbf{m}}$ the virtual transform of f at $p_{r-1}(K)$ relative to $\mathbf{m}$ is

$$\breve{f} = \sum_{i,j \geq 0} A^{r-1}_{ij}\left(f, \lambda \circ \psi_{r-1}(K)\right) \left(\breve{x} \circ i_{\breve{\breve{K}}}\right)^i \left(\breve{y} \circ i_{\breve{\breve{K}}}\right)^j .$$

This virtual transform must have multiplicity at least m_{r-1} at $p_{r-1}(K)$, therefore

$$A^{r-1}_{ij}(f, \lambda \circ \psi_{r-1}(K)) = 0 \ \forall i, j, \ i + j < m_{r-1} \ ,$$

and the virtual transform transform of $\breve{f}$ in $S_{r-1}(K)$ is given locally by

$$\sum_{i+j \geq m_{r-1}} A^{r-1}_{ij}(f, \lambda \circ \psi_{r-1}(K)) \left(\breve{x} \circ i_{\breve{\breve{K}}}\right)^{i+j-m_{r-1}} \left(\frac{\breve{y}}{\breve{x}} \circ i_{\breve{\breve{K}}}\right)^j = 0,$$

which because of the commutativity $i_{\breve{\breve{K}}} \circ \sigma_{r-1} = \pi_{r-1} \circ i_{\breve{K}}$ can be written in terms of the local parameters $x \circ i_{\breve{K}}, y \circ i_{\breve{K}}$:

$$\sum_{i+j \geq m_{r-1}} A^{r-1}_{ij}(f, \lambda \circ \psi_{r-1}(K)) \left(x \circ i_{\breve{K}}\right)^{i+j-m_{r-1}} \left(y \circ i_{\breve{K}} + \lambda_r(K)\right)^j =$$

$$= \sum_{i,j \geq m_{r-1}} \sum_{\ell=0}^{j} A^{r-1}_{ij}(f, \lambda \circ \psi_{r-1}(K)) \binom{j}{n} \lambda_r(K)^{j-\ell} \left(x \circ i_{\breve{K}}\right)^{i+j-m_{r-1}} \left(y \circ i_{\breve{K}}\right)^{\ell} \ .$$

This allows us to define the polynomials

$$A^r_{k\ell}(f,u_1,\ldots,u_{r-1}) = \sum_{i+j=k+m_{r-1}} A^{r-1}_{ij}(f,u_1,\ldots,u_{r-2})\binom{j}{n}u_{r-1}^{j-\ell}\ ,$$

after which the claim is satisfied. □

Proposition 3.2. *$H_{\mathbf{m}}$ is Zariski–closed in $\mathcal{O}\times Y_{r-1}$.*

Proof. By induction on r. The case $r=1$ is clear, because then $H_{K,\mathbf{m}} = \mathfrak{m}^{m_1}$ is the closed subset of $\mathcal{O} = \mathcal{O}\times Y_0$ determined by the ideal $I = (X_{ij})_{i+j<m_1}$. Suppose now $r>1$ and the claim is true for $H_{\breve{\mathbf{m}}} \subset \mathcal{O}\times Y_{r-2}$. Let $K_0 \in Y_{r-1}$ be a cluster and $U \subset Y_{r-1}$ the open neighbourhood given by lemma 3.1. It will be enough to see that $H_{U,\mathbf{m}} := H_{\mathbf{m}} \cap \pi^{-1}_{Y_{r-1}}(U)$ is closed in $\mathcal{O}\times U$, because Y_{r-1} can be covered by a finite number of such open neighbourhoods. Define

$$H'_{U,\mathbf{m}} = \{(f,K)\in \mathcal{O}\times U \mid (f,p_r(K)) \in H_{\breve{\mathbf{m}}}\}\ .$$

Because of the induction hypothesis and the fact that

$$id_{\mathcal{O}}\times p_r : \mathcal{O}\times Y_{r-1} \longrightarrow \mathcal{O}\times Y_r$$

is continuous, $H'_{U,\mathbf{m}}$ is closed in $\mathcal{O}\times U$. Moreover $(f,K)\in H_{U,\mathbf{m}}$ if and only if $(f,K)\in H'_{U,\mathbf{m}}$ and the virtual transform of $f=0$ at $p_r(K)$ has multiplicity at least m_r, that is, by lemma 3.1,

$$A^r_{ij}(f,\lambda(K)) = 0\ \forall i,j,\ i+j<m_r\ .$$

These equations define a closed subset $V(A^r_{ij})_{i+j<m_r} \subset \mathcal{O}\times U$, therefore $H_{U,\mathbf{m}} = H'_{U,\mathbf{m}} \cap V(A^r_{ij})_{i+j<m_r}$ is closed in $\mathcal{O}\times U$. □

From this we obtain a number of corollaries. Let N be a positive integer such that $\dim(\mathcal{O}/H_{K,\mathbf{m}}) \le N$ for all $K\in Y_{r-1}$ (for example, $N = \sum_{i=1}^r m_i\,(m_i+1)/2$). Then $H_{K,\mathbf{m}} \supset \mathfrak{m}^N$ for all K, and we can define

$$\bar{H}_{\mathbf{m}} := \{(\bar{f},K)\in \frac{\mathcal{O}}{\mathfrak{m}^N}\times Y_{r-1} \mid f\in H_{K,\mathbf{m}}\}\,.$$

Corollary 3.3. *$\bar{H}_{\mathbf{m}}$ is Zariski–closed in $\left(\mathcal{O}/\mathfrak{m}^N\right)\times Y_{r-1}$.*

Proof. Let $K_0\in Y_{r-1}$ be a cluster and $U\subset Y_{r-1}$ the open neighbourhood given by lemma 3.1. As in lemma 3.2, it will be enough to see that $\bar{H}_{U,\mathbf{m}} := \bar{H}_{\mathbf{m}} \cap \pi^{-1}_{Y_{r-1}}(U)$ is closed in $(\mathcal{O}/\mathfrak{m}^N)\times U$. We also know from the proof of lemma 3.2 that $H_{U,\mathbf{m}}$ is defined by a finite number of polynomials A^k_{ij}, $k=1,\ldots,r$. Now for $\alpha+\beta\ge N$, $x^\alpha y^\beta \in H_{K,\mathbf{m}}$ for all K, so the polynomial $A^k_{ij}(x^\alpha y^\beta,u_1,\ldots,u_{k-1}) \in k[u_1,\ldots,u_{k-1}]$ must be identically zero. This implies in fact that

$$A^k_{ij} \in k[X_{\alpha\beta}]_{\alpha+\beta<N}[u_1,u_2,\ldots,u_{k-1}] \subset R[u_1,u_2,\ldots,u_{k-1}]\ .$$

Now $k[X_{\alpha\beta}]_{\alpha+\beta<N}$ is the affine coordinate ring of $\mathcal{O}/\mathfrak{m}^N$, which is a k–vector space admiting the basis $\{\bar{x}^\alpha\bar{y}^\beta\}_{\alpha+\beta<N}$. Therefore the A^k_{ij} define a Zariski–closed subset in $\left(\mathcal{O}/\mathfrak{m}^N\right)\times U$, and it is immediate to see that this is in fact $\bar{H}_{U,\mathbf{m}}$. □

Corollary 3.4. *The function*

$$\begin{aligned} Y_{r-1} &\longrightarrow \mathbb{Z}_{\geq 0} \\ K &\longmapsto \dim \frac{\mathcal{O}}{H_{K,\mathbf{m}}} = \text{length } Z_{K,\mathbf{m}} \end{aligned}$$

is lower semicontinuous.

Proof. Choose N such that $\dim(\mathcal{O}/H_{K,\mathbf{m}}) \leq N$ for all $K \in Y_{r-1}$. Then

$$\dim \frac{\mathcal{O}}{H_{K,\mathbf{m}}} = \dim \frac{\mathcal{O}/\mathfrak{m}^N}{H_{K,\mathbf{m}}/\mathfrak{m}^N} = \frac{N(N+1)}{2} - \dim \left(\bar{H}_{\mathbf{m}} \times_{Y_{r-1}} \{K\}\right)$$

and the claim follows because the dimension of the fibers of the (non flat) family $\bar{H}_{\mathbf{m}} \to Y_{r-1}$ is upper semicontinuous. □

Remark that in general there is no ideal sheaf $\mathcal{I}$ on $S \times Y_{r-1}$ with $\mathcal{I} \otimes k(K) = H_{K,\mathbf{m}}$ for all K; in other words, all $Z_{K,\mathbf{m}}$ do not form a family. Indeed, the length of the members of a family of zero–dimensional schemes is upper semicontinuous, against corollary 3.4. As the simple example $\mathbf{m} = (2,2,2)$ shows, this length is not always constant, and in fact the systems of multiplicities for which it is constant are quite exceptional. However, we will see that restricting the set of clusters under consideration so that the length of the corresponding schemes remains constant, it is possible to construct flat families of cluster schemes.

Given a subvariety $W \subset Y_{r-1}$, let $W_{\mathbf{m}} \subset W$ be the open subset where $\dim(\mathcal{O}/H_{K,\mathbf{m}})$ is maximal, and denote this dimension by $N = N(\mathbf{m}, W)$. The restriction

$$\bar{H}_{\mathbf{m}}|_{W_{\mathbf{m}}} \subset \frac{\mathcal{O}}{\mathfrak{m}^N} \times W_{\mathbf{m}}$$

is thus a family of vector subspaces of $\mathcal{O}/\mathfrak{m}^N$ of codimension N, so it defines a morphism $W_{\mathbf{m}} \to \mathbb{G}$ to the Grassmannian $\mathbb{G} = \mathbb{G}_N\left(\mathcal{O}/\mathfrak{m}^N\right)$. In $\mathbb{G}$, the codimension N vector spaces which are ideals of $\left(\mathcal{O}/\mathfrak{m}^N\right)$ constitute a closed subscheme, which is identified to the Hilbert scheme $\text{Hilb}_p^N S$ of length N subschemes of S supported at p (cf. [13], [25]). As $H_{K,\mathbf{m}}$ is by definition an ideal for every K, we finally obtain a morphism

$$\begin{aligned} W_{\mathbf{m}} &\xrightarrow{\varphi_{\mathbf{m}}} \text{Hilb}^N S \\ K &\longmapsto Z_{K,\mathbf{m}} \end{aligned}$$

That is, the schemes $Z_{K,\mathbf{m}}$ form a flat family with parameter space $W_{\mathbf{m}}$. In the particular case that $W = C$ is a smooth curve, the morphism $\varphi_{\mathbf{m}}$ can be extended to all of C, by [22, III, 9.8], but then it is not true that $\varphi_{\mathbf{m}}(K) = Z_{K,\mathbf{m}}$ for those $K \in C \setminus C_{\mathbf{m}}$. In fact, $\varphi_{\mathbf{m}}(K)$ depends not only on K and $\mathbf{m}$ but also on C. Nevertheless, we have the following

Corollary 3.5. *If $C \subset Y_{r-1}$ is a smooth curve and $K \in C$ then $Z_{K,\mathbf{m}} \subseteq \varphi_{\mathbf{m}}(K)$.*

Proof. The morphism $\varphi_{\mathbf{m}} : C \to \text{Hilb}_p^N S \subset \mathbb{G}$ defines a closed subset $\Phi_{\mathbf{m}} \subset (\mathcal{O}/\mathfrak{m}^N) \times C$, and we have to see that $\Phi_{\mathbf{m}} \subset \bar{H}_{\mathbf{m}}|_C$. As $\bar{H}_{\mathbf{m}}$ is closed and

$$\Phi_{\mathbf{m}}|_{C_{\mathbf{m}}} = \bar{H}_{\mathbf{m}}|_{C_{\mathbf{m}}} \subset \frac{\mathcal{O}}{\mathfrak{m}^N} \times C_{\mathbf{m}}$$

it will be enough to see that $\Phi_{\mathbf{m}}$ is the closure of $\Phi_{\mathbf{m}}|_{C_{\mathbf{m}}}$.

Take $(\bar{f}, K_0) \in \Phi_{\mathbf{m}}$, with $K_0 \in C \setminus C_{\mathbf{m}}$. The family determined by $\varphi_{\mathbf{m}}$ corresponds to an ideal sheaf $\mathcal{I} \subset (\mathcal{O}/\mathfrak{m}^N) \otimes \mathcal{O}_C$. By definition, $\bar{f} \in \mathcal{I} \otimes_{\mathcal{O}_C} k(K_0)$, so there is an open neigbourhood U of K_0 in C and a section

$$\bar{f}_U \in \Gamma\left(\operatorname{Spec} \frac{\mathcal{O}}{\mathfrak{m}^N} \times U, \mathcal{I}\right)$$

with $\bar{f} = \bar{f}_U(K_0)$. As U is open, we may assume $U \setminus \{K_0\} \subset C_{\mathbf{m}}$, therefore $\forall K \in U \setminus \{K_0\}$, $\bar{f}_U(K) \in \mathcal{I} \otimes k(K) = \bar{H}_{K,\mathbf{m}}$. So we have a morphism

$$\begin{aligned} U &\longrightarrow \Phi_{\mathbf{m}} \\ K &\mapsto (\bar{f}_U(K), K) \end{aligned}$$

whose image contains $(\bar{f}, K_0)$ and has every other point in $\Phi_{\mathbf{m}}|_{C_{\mathbf{m}}}$, therefore $(\bar{f}, K_0)$ lies in the closure of $\Phi_{\mathbf{m}}|_{C_{\mathbf{m}}}$. As this holds for every $(\bar{f}, K_0) \in \Phi_{\mathbf{m}}$ and $\Phi_{\mathbf{m}}$ is closed, the claim is proved. □

Next lemma shows an example of a flat family of cluster schemes which will be useful later on.

Lemma 3.6. *Let $(K, \mathbf{m})$ be a consistent weighted cluster of r points and $E = E_r \subset S_K$ the exceptional divisor of blowing up the last point. Consider the system of multiplicities $\mathbf{m}_1 = (m_1, m_2, \ldots, m_r, 1)$ and for every $q \in E$, $K_q = (p_1, p_2, \ldots, p_r, q)$. Then the schemes $Z_{K_q,\mathbf{m}_1}$ form a flat family.*

Proof. It is clear that $\operatorname{length} Z_{K_q,\mathbf{m}_1} \leq Z_{K,\mathbf{m}} + 1 \ \forall q$. On the other hand, as $(K, \mathbf{m})$ is consistent, there are curves going sharply through $(K, \mathbf{m})$ which miss q (cf. [5, 4]), so $\operatorname{length} Z_{K_q,\mathbf{m}_1} = Z_{K,\mathbf{m}} + 1 \ \forall q$. Abusing slightly notations, we will call $E = i_K(E) \subset Y_r$; as the length of $Z_{K_q,\mathbf{m}_1}$ does not depend on q we have $E_{\mathbf{m}_1} = E$ and the schemes $Z_{K_q,\mathbf{m}_1}$ form a flat family.

□

Recall that $P_s \subset Y_{r-1}$ is the irreducible subvariety containing the unibranched clusters K with $p_s(K)$ proximate to p_1, and $U_s \subset P_s$ is the open dense subset which contains all unibranched clusters based at p_1 with $p_i(K)$ proximate to $p_{i-1}(K)$ for all $i > 1$ and to $p_1(K)$ if $s \geq i > 1$, and no other proximity relations. Assume from now that the system of multiplicities we are dealing with is of the form $\mathbf{m} = (m, 2^i, 1^j)$. We will denote by $\mathcal{V}(s, m, i, j) = \overline{\varphi_{\mathbf{m}}(U_s)}$ the closure of the subscheme of the Hilbert scheme that parametrizes the unibranched cluster schemes $Z_{K,(m,2^i,1^j)}$ with $K \in U_s$. Note that this does not depend on r, as far as $r \geq i + j + 1$. Note also that $U_s \subset (P_s)_{\mathbf{m}}$ because U_s is open and dense in P_s and $\operatorname{length} Z_{K,\mathbf{m}}$ is constant on U_s. Therefore $\varphi_{\mathbf{m}}((P_s)_{\mathbf{m}}) \subset \overline{\varphi_{\mathbf{m}}(U_s)}$.

Proposition 3.7. *Suppose that $\mathbf{m} = (m, 2^i, 1^j)$ is consistent in U_s and $(P_s)_{\mathbf{m}}$ does not contain U_{s+1}. Then*

1. *$m = 2s - 2$ and $i \geq s$,*
2. *if $i + j \geq s^2 - 3s + 1$ then $\mathcal{V}(s, m, i, j)$ contains $\mathcal{V}(s + 1, m + 1, i - s + 1, j + s - 2)$,*
3. *$(m + 1, 2^{i-s+1}, 1^{j+s-2})$ is consistent in U_{s+1} if and only if $i \leq 2s - 2$. If it is not, then it is equivalent to $(m + 2, 2^{i-2s+1}, 1^{j+2s-2})$.*



Figure 2: Enriques diagrams illustrating proposition 3.7. By moving the free points, $Z_{K,\mathbf{m}}$ specializes to $Z_{K',\mathbf{m}'}$.

Proof. 1. $\mathbf{m}$ is consistent in U_s, but not consistent in U_{s+1}, otherwise $U_{s+1} \subset (P_s)_{\mathbf{m}}$. Therefore the proximity inequality at p_1 must be satisfied for $K \in U_s$ but not for $K \in U_{s+1}$. This means that

$$\begin{cases} 2\,s > m \geq 2\,s-2 & \text{if } i \geq s, \\ s+i > m \geq s+i-1 & \text{if } i < s. \end{cases}$$

Therefore, if $i < s$ we must have $m = s+i-1$, and for any cluster $K \in U_{s+1}$ we have $\delta(\mathbf{m}) = (m+1, 1^{2i+j-s})$. So

$$\text{length}\, Z_{K,\mathbf{m}} = \text{length}\, Z_{K,\delta(\mathbf{m})} =$$
$$= \frac{(m+1)(m+2)}{2} + 2\,i + j - s = \frac{m\,(m+1)}{2} + 3\,i + j$$

for all $K \in U_{s+1}$, so $U_{s+1} \subset (P_s)_{\mathbf{m}}$ against the hypotheses. We conclude that $i \geq s$, and it only remains to be seen that the case $m = 2\,s-1$ is not possible. But in this case $\delta(\mathbf{m}) = (m+1, 2^{i-s}, 1^{j+s})$ in U_{s+1}, so again

$$\text{length}\, Z_{K,\mathbf{m}} = \text{length}\, Z_{K,\delta(\mathbf{m})} =$$
$$= \frac{(m+1)(m+2)}{2} + 3\,(i-s) + j + s = \frac{m\,(m+1)}{2} + 3\,i + j$$

for all $K \in U_{s+1}$ and $U_{s+1} \subset (P_s)_{\mathbf{m}}$.

2. We are given two systems of multiplicities

$$\mathbf{m} = (2\,s-2, 2^i, 1^j)$$
$$\mathbf{m}' = (2\,s-1, 2^{i-s+1}, 1^{j+s-2})$$

with $i > s$ and $i+j+1 \geq (s-1)(s-2)$. We have to see that for any cluster $K_0 \in U_{s+1}$ there is a deformation of $Z_{K_0,\mathbf{m}'}$ whose general member

is of the form $Z_{K,\mathbf{m}}$ with $K \in U_s$. For that, let C be a smooth irreducible curve $K_0 \in C \subset P_s$ with $C \cap U_s \neq \emptyset$ and consider the morphism

$$C \xrightarrow{\varphi_{\mathbf{m}}} \mathrm{Hilb}^N S\,.$$

Consider also an auxiliary system of multiplicities and the associated morphism

$$\mathbf{m}_+ = (2s-1, 2^{i-s+1}, 1^{j+s-1})$$
$$C \xrightarrow{\varphi_{\mathbf{m}_+}} \mathrm{Hilb}^{N+1} S\,.$$

For $K \in C \cap U_s$, $Z_{K,\mathbf{m}} \subsetneq Z_{K,\mathbf{m}_+}$. Therefore $\varphi_{\mathbf{m}}(K) \subsetneq \varphi_{\mathbf{m}_+}(K)$ for all $K \in C$. Moreover, part 1 of the proposition shows that $K_0 \in C_{\mathbf{m}_+}$, so $\varphi_{\mathbf{m}_+}(K_0) = Z_{K_0,\mathbf{m}_+}$. On the other hand, $(K_0, \mathbf{m})$ does not satisfy the proximity inequality at p_1, and unloading multiplicity on this point gives

$$\mathbf{m}_- = (2\,s-1, 2^{i-s}, 1^{j+s})\,.$$

Note that $\mathbf{m}_-$ may be non–consistent, but still $Z_{K_0,\mathbf{m}} = Z_{K_0,\mathbf{m}_-}$ and $\operatorname{length} Z_{K_0,\mathbf{m}} < \operatorname{length} \varphi_{\mathbf{m}}(K_0)$. All together, we have

$$Z_{K_0,\mathbf{m}_-} = Z_{K_0,\mathbf{m}} \subsetneq \varphi_{\mathbf{m}}(K_0) \subsetneq \varphi_{\mathbf{m}_+}(K_0) = Z_{K_0,\mathbf{m}_+}$$

and we are in the conditions of lemma 2.1. Therefore there is a point q in the exceptional divisor $E = E_r$ of blowing up p_r such that $\varphi_{\mathbf{m}}(K_0) = Z_{K_q,\mathbf{m}_0}$, with

$$\mathbf{m}_0 = (2\,s-1, 2^{i-s}, 1^{j+s+1})\,.$$

Let $q_0 \in E$ be the only point proximate to p_{r-1}. If $q = q_0$ then unloading gives $Z_{K_q,\mathbf{m}_0} = Z_{K_0,\mathbf{m}'}$ so $\varphi_{\mathbf{m}}$ is the family we are looking for. If q is not proximate to p_{r-1} then we only know that $Z_{K_q,\mathbf{m}_0} \in \mathcal{V}(s,m,i,j)$. As $i+j+1 \geq (s-1)(s-2)$, though, by lemma 2.4 we can say that for any free point $q \in E$, $Z_{K_q,\mathbf{m}_0} \in \mathcal{V}(s,m,i,j)$. As the free points are dense in E, this implies $\varphi_{\mathbf{m}_0}(E) \subset \mathcal{V}(s,m,i,j)$. Now it is enough to see that $q_0 \in E_{\mathbf{m}_0}$, because then $Z_{K_0,\mathbf{m}'} = Z_{K_{q_0},\mathbf{m}_0} = \varphi_{\mathbf{m}_0}(q_0) \in \mathcal{V}(s,m,i,j)$. If $(K_q, \mathbf{m}_0)$ is consistent for $q \in E$ free, then $q_0 \in E_{\mathbf{m}_0}$ because of lemma 3.6; if it is not, then the equivalent consistent system obtained by unloading is $\delta_{K_q}(\mathbf{m}_0) = (m+2, 2^{i-2s}, 1^{j+2s+1})$, whereas for the cluster K_{q_0} the equivalent consistent system is $\delta_{K_{q_0}}(\mathbf{m}_0) = (m+2, 2^{i-2s+1}, 1^{j+2s-2})$. This implies that $\operatorname{length} Z_{K_q,\mathbf{m}_0} = N(\mathbf{m}_0, E)$, so again $q_0 \in E_{\mathbf{m}_0}$.

3. Follows from an easy unloading calculation.

□

Corollary 3.8. *Suppose that $\mathbf{m} = (m, 2^i, 1^j)$ is consistent in U_s and $(P_s)_{\mathbf{m}}$ does not contain U_{s+1}, and suppose that $j \geq s^2 - 5\,s + 2$. For every $k \in \mathbb{Z}_{\geq 0}$ we define*

$$m_k = m + 2\,k$$
$$i_k = i - k\,(k + 2\,s - 2)$$
$$j_k = j + k\,(k + 2\,s - 3)\,.$$

If either $k = 0$ or $i_{k-1} > 2\,(s+k-1) - 2$ then $\mathcal{V}(s+k, m_k, i_k, j_k) \subset \mathcal{V}(s,m,i,j)$.

Proof. We will proceed by induction on k. For $k = 0$, there is nothing to prove. For $k > 0$, we have either $k - 1 = 0$ or

$$i_{k-2} > i_{k-1} > 2(s + k - 1) - 2 > 2(s + k - 2) - 2$$

so we can apply the induction hypothesis and

$$\mathcal{V}(s + k - 1, m_{k-1}, i_{k-1}, j_{k-1}) \subset \mathcal{V}(s, m, i, j).$$

On the other hand, a straightforward computation shows that $j_{k-1} \geq (s + k - 1)^2 - 5(s + k - 1) + 2$, therefore $i + j \geq (s + k - 1)^2 - 3(s + k - 1) + 1$. In these contitions, proposition 3.7 tells us that

$$\mathcal{V}(s + k, m_k, i_k, j_k) \subset \mathcal{V}(s + k - 1, m_{k-1}, i_{k-1}, j_{k-1}),$$

thus finishing the proof. □

4 Degree of singular plane curves

In this section we work on a projective irreducible smooth surface S, mainly $S = \mathbb{P}^2$. We are interested in linear systems of curves which contain a zero dimensional scheme composed of unibranched cluster schemes. These linear systems can be specialized to linear systems of curves through a unibranched cluster (supported at a single point) using the technique developped in [34]. We will not repeat the whole treatment here, but only show the parameter space (a variety of clusters) to which we apply the semicontinuity theorem.

For convenience, we fix a point p in S. Given a sequence of integers $\mathbf{r} = (r_1, r_2, \ldots, r_\rho)$ with $r_1 < r_2 < \ldots < r_\rho$ we define varieties $X_i(\mathbf{r})$ and $Y_i(\mathbf{r})$ in an analogous way to X_i and Y_i. Let $Y_{-1}(\mathbf{r}) = \operatorname{Spec} k$, $X_0(\mathbf{r}) = S$, $Y_0(\mathbf{r}) = \{p\}$, and for $i > 0$ let

$$X_i(\mathbf{r}) \xrightarrow{b_i} Y_{i-1}(\mathbf{r}) \times_{Y_{i-2}(\mathbf{r})} X_{i-1}(\mathbf{r})$$

be the blowing-up along $\Delta(Y_{i-1}(\mathbf{r}))$,

$$Y_i(\mathbf{r}) = \begin{cases} \text{exc. divisor of the blowing-up} & \text{if } i \notin \{r_1, r_2, \ldots, r_\rho\} \\ X_i(\mathbf{r}) & \text{if } i \in \{r_1, r_2, \ldots, r_\rho\} \end{cases}$$

and $\psi_i(\mathbf{r}) = \pi_{Y_{i-1}(\mathbf{r})} \circ b_i : X_i(\mathbf{r}) \to Y_{i-1}(\mathbf{r})$ is smooth of relative dimension 2. We write also $X(\mathbf{r}) = X_{r_\rho}(\mathbf{r})$ and $Y(\mathbf{r}) = Y_{r_\rho - 1}(\mathbf{r})$. All these varieties are projective, irreducible and smooth; $Y(\mathbf{r})$ can be identified with the set of all clusters of r_ρ points of S, such that $p_1 = p$ and p_{i+1} is proximate to p_i for all $i \notin \mathbf{r}$. A general cluster $K \in Y(\mathbf{r})$ is the union of ρ unibranched clusters of $r_1, r_2 - r_1, \ldots,$ and $r_\rho - r_{\rho-1}$ points.

Lemma 4.1. *For any sequence of integers $\mathbf{r} = (r_1, r_2, \ldots, r_\rho)$, X_i and Y_i are closed subvarieties of $X_i(\mathbf{r})$ and $Y_i(\mathbf{r})$ respectively, and the morphism $\psi_i : X_i \to Y_{i-1}$ of section 2 is the restriction of $\psi_i(\mathbf{r}) : X_i(\mathbf{r}) \to Y_{i-1}(\mathbf{r})$ for all i.*

Proof. By induction on i. The cases $i \leq 1$ are clear, so assume $i > 1$ and the claim to be true for X_{i-1} and X_{i-2}. This means that $X_{i-1} \subset X_{i-1}(\mathbf{r})$, $Y_{i-1} \subset Y_{i-1}(\mathbf{r})$ are closed subvarieties and the morphism $\psi_{i-1} : X_{i-1} \to Y_{i-2}$

is the restriction of $\psi_{i-1}(\mathbf{r}) : X_{i-1}(\mathbf{r}) \to Y_{i-2}(\mathbf{r})$. So we have also a closed subvariety

$$Y_{i-1} \times_{Y_{i-2}} X_{i-1} \subset Y_{i-1}(\mathbf{r}) \times_{Y_{i-2}(\mathbf{r})} X_{i-1}(\mathbf{r})$$

and

$$\Delta(Y_{i-1}(\mathbf{r})) \cap Y_{i-1} \times_{Y_{i-2}} X_{i-1} = \Delta(Y_{i-1})$$

so by the definitions, $X_i \subset X_{i-1}(\mathbf{r})$ is the strict transform of $Y_{i-1} \times_{Y_{i-2}} X_{i-1}$ under the blowing-up b_i (cf. [22, II.7.15]), Y_i is a subvariety of the exceptional divisor, hence of $Y_{i-1}(\mathbf{r})$, and $\psi_i : X_i \to Y_{i-1}$ is the restriction of $\psi_i : X_i(\mathbf{r}) \to Y_{i-1}(\mathbf{r})$. □

Given a cluster $K \in Y(\mathbf{r})$, a system of multiplicities $\mathbf{m}$ and a divisor D, we will denote by $\mathcal{L}_D(K, \mathbf{m})$ the linear system of effective divisors linearly equivalent to D which go through $(K, \mathbf{m})$. As seen in [34] the function

$$\begin{aligned} Y(\mathbf{r}) &\longrightarrow \mathbb{Z}_{\geq -1} \\ K &\longmapsto \ell_D(K, \mathbf{m}) := \dim \mathcal{L}_D(K, \mathbf{m}) \end{aligned}$$

is upper semicontinuous. We consider the dimension of the empty linear system to be -1. Because of lemma 4.1, we can bound $\ell_D(K, \mathbf{m})$ with K general in $Y(\mathbf{r})$ by $\ell_D(K, \mathbf{m})$ with K in $Y_{r_\rho - 1}$. For any subset $W \subset Y(\mathbf{r})$ we will also write $\ell_D(W, \mathbf{m}) = \inf\{\ell_D(K, \mathbf{m}) \mid K \in W\}$.

In the case $S = \mathbb{P}^2$, we can choose $D = d \cdot L$, with L a line and $d \in \mathbb{Z}_{>0}$. We will then denote $\mathcal{L}_d(K, \mathbf{m}) = \mathcal{L}_D(K, \mathbf{m})$ the linear system of curves of degree d going through $(K, \mathbf{m})$ and $\ell_d(K, \mathbf{m}) = \ell_D(K, \mathbf{m})$ its dimension. We say that a zero–dimensional scheme Z has level d when $\operatorname{length} Z = (d+1)(d+2)/2$. For a level d scheme in the plane $\mathbb{P}^2$ it is equivalent to have maximal rank or to have maximal rank in degree d. Furthermore, an arbitrary zero–dimensional subscheme $Z \subset \mathbb{P}^2$ has maximal rank if and only if there exist maximal rank schemes Z_d and Z_{d+1} of level d and $d+1$ respectively such that $Z_d \subset Z \subset Z_{d+1}$ (see [23, 2.2.2]).

Lemma 4.2. *For every cluster scheme $Z_{K,\mathbf{m}}$ where $K \in U_s$ and $\mathbf{m} = (m, 2^i, 1^j)$ is consistent in U_s there exist systems of multiplicities $\mathbf{m}_- = (m, 2^{i_-}, 1^{j_-})$ and $\mathbf{m}_+ = (m, 2^{i_+}, 1^{j_+})$ consistent in U_s such that $Z_{K,\mathbf{m}_-} \subset Z_{K,\mathbf{m}} \subset Z_{K,\mathbf{m}_+}$, $Z_{K,\mathbf{m}_-}$ has level d and $Z_{K,\mathbf{m}_+}$ has level $d+1$. Furthermore, if $4j \geq m^2 - 4m - 6$ and $3i + j \geq 2m + 3$ then $\mathbf{m}_+$ and $\mathbf{m}_-$ can be chosen such that $4j_+ \geq m^2 - 4m - 6$, $4j_- \geq m^2 - 4m - 6$ or $i_- = 0$, and $d > m$.*

Proof. Write

$$\frac{m(m+1)}{2} + 3\,i + j = \frac{(d+1)(d+2)}{2} + \epsilon$$

with $0 \leq \epsilon \leq d+1$. It is enough to define $\mathbf{m}_+ = (m, 2^i, 1^{j+d+2-\epsilon})$ and

$$\mathbf{m}_- = \begin{cases} (m, 2^{i-\epsilon}, 1^{j+2\epsilon}) & \text{if } i \geq \epsilon \\ (m, 1^{j+3i-\epsilon}) & \text{if } i \leq \epsilon \end{cases}$$

□

Before we prove our maximal rank theorem we need some lemmas on level d cluster schemes. Assume for a while that $\mathbf{m} = (m, 2^i, 1^j)$ and $d \in \mathbb{Z}$ are such that

$$N(\mathbf{m}) = \frac{m\,(m+1)}{2} + 3\,i + j = \frac{(d+1)\,(d+2)}{2}\,.$$

Lemma 4.3. *If $i = 0$ and $s \leq m+1$ then $\ell_d(U_s, \mathbf{m}) = -1$, with $U_s \subset Y_{r_\rho - 1}$ as defined in section 3.*

Proof. It is clear that $d \geq m-1$; we will prove the claim by induction on $d-m$. For $d = m-1$ the result is obvious. For $d \geq m$ we have $j \geq m+1$. By semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_{m+1}. Now for $K \in U_{m+1}$, unloading gives $Z_{K,\mathbf{m}} = Z_{K,(m+1,1^{j-m-1})}$, and the result follows from the induction hypothesis. □

Lemma 4.4. *If $2\,i \leq m$, $2\,i+j > m$ and $\mathbf{m}$ is consistent in U_s then $\ell_d(U_s, \mathbf{m}) = -1$.*

Proof. By semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_{m+1-i}. But for $K \in U_{m+1-i}$, unloading gives $Z_{K,\mathbf{m}} = Z_{K,(m+1,1^{2i+j-m-1})}$, and the result follows from lemma 4.3. □

Proposition 4.5. *If $d = m+1$, $m \geq 2i-2$ and $\mathbf{m}$ is consistent in U_s then $\ell_d(U_s, \mathbf{m}) = -1$.*

Proof. The proof runs by induction on i, the case $i = 0$ being clear from lemma 4.3. Assume $i \geq 1$. For K general in U_s, consider the line L passing through p in the direction of p_2. If $s = 2$ we can assume (K being general) that p_3 does not lie on L; for $s > 2$ this is automatic. For every curve $C \in \mathcal{L}_d(K, \mathbf{m})$

$$L \cdot C \geq m+2 = d+1 > \deg C,$$

therefore L is a component of C, so it is a fixed part of $\mathcal{L}_d(K, \mathbf{m})$. The residual linear system is

$$\mathcal{L}_d(K, \mathbf{m}) - L = \mathcal{L}_{d-1}(K, \mathbf{m}')$$

with $\mathbf{m}' = (m-1, 1, 2^{i-1}, 1^j)$ because of lemma 2.3. As this is true for general $K \in U_s$ we have now $\ell_d(U_s, \mathbf{m}) = \ell_{d-1}(U_s, \mathbf{m}')$. But $\mathbf{m}'$ is not consistent in U_s. The equivalent consistent system is

$$\delta(\mathbf{m}') = \begin{cases} (m, 1^{j+i}) & \text{if } m = 2\,(s-1) = 2\,(i-1) \\ (m-1, 2^{i-1}, 1^{j+1}) & \text{in any other case.} \end{cases}$$

Observe that $N(\delta(\mathbf{m})) = d(d+1)/2$. Finally $\ell_d(U_s, \mathbf{m}) = \ell_{d-1}(U_s, \delta(\mathbf{m}')) = -1$ because of lemma 4.3 in the first case and because of the induction hypothesis in the second. □

Corollary 4.6. *If $2i \leq m$, $d > m$ and $\mathbf{m}$ is consistent in U_s then $\ell_d(U_s, \mathbf{m}) = -1$.*

Proof. By lemma 4.4 we may assume $2i + j \leq m$. But then

$$N(\mathbf{m}) = \frac{m(m+1)}{2} + 3\,i + j \leq \frac{m(m+1)}{2} + m + \frac{m}{2} < \frac{(m+3)(m+4)}{2}$$

so $d < m + 2$, and proposition 4.5 concludes. □

Lemma 4.7. *If $i \leq m$, $d > m$, $4\,(i+j) \geq m^2 - 2\,m - 4$ and $\mathbf{m}$ is consistent in U_s then $\ell_d(U_s, \mathbf{m}) = -1$.*

Proof. By corollary 4.6 we may assume $2i > m$. We will distinguish two cases according to the parity of m.

m even Write $m = 2\,t - 2$. As $\mathbf{m}$ is consistent in U_s, and $2i > m$, we must have $t \geq s$, so by semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_t. It is clear that $(P_t)_{\mathbf{m}}$ does not contain U_{t+1}, and $4\,(i+j) \geq 4\,(t^2 - 3\,t + 1)$, therefore by proposition 3.7

$$\mathcal{V}(t+1, m+1, i-t+1, j+t-2) \subset \mathcal{V}(t, m, i, j)\ .$$

By semicontinuity applied to the tautological flat family on $\mathcal{V}(t, m, i, j) \subset \mathrm{Hilb}^N(S)$ then, it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}'}$ with K general in U_{t+1} and $\mathbf{m}' = (m+1, 2^{i-t+1}, 1^{j+t-2})$. But then $2(i-t+1) = 2i - m \leq m$ and still $2(i-t+1) + j + t - 2 > m$ so the result follows by lemma 4.4.

m odd Write $m = 2\,t - 1$. By semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_{t+1}. But for $K \in U_{t+1}$, unloading gives $Z_{K,\mathbf{m}} = Z_{K,(m+1,2^{i-t},1^{j+t})}$, and again $2(i-t) = 2i - m - 1 \leq m$ and $2(i-t) + j + t > m$ so the result follows by lemma 4.4.

□

With this knowledge of level d cluster schemes we are in a position to attack the general case. Let now ρ, s be positive integers, and suppose we have systems of multiplicities $\mathbf{m}_1, \mathbf{m}_2, \ldots, \mathbf{m}_\rho$ with

$$\begin{aligned} \mathbf{m}_1 &= (m, 2^{i_1}, 1^{j_1}) \\ \mathbf{m}_k &= (2^{i_k}, 1^{j_k}) \quad k = 2, 3, \ldots, \rho. \end{aligned}$$

Suppose furthermore that $m \geq \min(s + \sum i_k, 2s)$. Consider a cluster scheme $Z_1 = Z(K, \mathbf{m}_1)$ with $K \in U_s$ and $\rho - 1$ unibranched cluster schemes $Z_2, Z_3, \ldots$, Z_ρ supported at different points of $\mathbb{P}^2$, whose defining clusters $K_2, K_3, \ldots, K_\rho$ have no satellite points. The scheme $Z = Z_1 \cup Z_2 \cup \cdots \cup Z_\rho$ has length

$$N = \frac{m\,(m+1)}{2} + 3 \sum_{k=0}^{\rho} i_k + \sum_{k=0}^{\rho} j_k\,.$$

Theorem 4.8. *If the position of the points of the ρ clusters $K_2, K_3, \ldots, K_\rho$ is general, $3 \sum i_k + \sum j_k \geq 2m + 3$, and $4 \sum j \geq m^2 - 4\,m - 6$, then Z has maximal rank, except in the cases*

- $m = 2$, $\sum i_k = 4$, $j_k = 0\ \forall k$,
- $m = 4$, $\sum i_k = 6$, $j_k = 0\ \forall k$.

Proof. Consider $\mathbf{r} = (r_1, r_2, \dots, r_\rho)$. Let $P_s(\mathbf{r})$ be the closed variety of $Y(\mathbf{r})$ where $p_2, p_3, \dots, p_s$ are proximate to p. It is easy to see that $P_s(\mathbf{r})$ is an irreducible variety and $P_s = P_s(\mathbf{r}) \cap Y_{r_\rho - 1}$. Then the claim may be equivalently stated as

$$\ell_d(P_s(\mathbf{r}), \mathbf{m}_0) = \max\left(-1, \frac{(d+1)(d+2)}{2} - N\right)\ \forall d$$

where $\mathbf{m}_0 = (m, 2^{i_1}, 1^{j_1}, 2^{i_2}, 1^{j_2}, \dots, 2^{i_\rho}, 1^{j_\rho})$. By semicontinuity it is enough to see that $\ell_d(P_s, \mathbf{m}_0) = \max(-1, (d+1)(d+2)/2 - N) \forall d$ and unloading gives that $\mathbf{m} = (m, 2^{\sum i_k}, 1^{\sum j_k})$ is equivalent to $\mathbf{m}_0$ in U_s. Therefore

$$\ell_d(P_s(\mathbf{r}), \mathbf{m}_0) \leq \ell_d(U_s, \mathbf{m}_0) = \ell_d(U_s, \mathbf{m})\ \forall d\ .$$

It only remains to be seen that for a general cluster $K \in U_s$, $Z_{K,\mathbf{m}}$ has maximal rank. Because of lemma 4.2 we can assume that $Z_{K,\mathbf{m}}$ has level $d > m$, and it is enough to see that $\ell_d(U_s, \mathbf{m}) = -1$.

Let $i = \sum i_k$, $j = \sum j_k$. The proof runs by induction on i. The case $i \leq m$ has already been settled in lemma 4.7, so suppose $i \geq m+1$. We will distinguish two cases according to the parity of m.

m even Write $m = 2\,t - 2$. As $\mathbf{m}$ is consistent in U_s and $i \geq m + 1$ we must have $t \geq s$, so by semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_t. It is clear that $(P_t)_{\mathbf{m}}$ does not contain U_{t+1}, and $4\,(i+j) \geq 4\,(t^2 - 3\,t + 1)$, therefore by proposition 3.7

$$\mathcal{V}(t+1, m+2, i-2t+1, j+2t-2) \subset \mathcal{V}(t, m, i, j)\ .$$

By semicontinuity applied to the tautological flat family on $\mathcal{V}(t, m, i, j) \subset \mathrm{Hilb}^N(S)$ then, it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}'}$ with K general in U_{t+1} and

$$\mathbf{m}' = (m', 2^{i'}, 1^{j'}) = (m+2, 2^{i-2t+1}, 1^{j+2t-2})\,.$$

An easy computation shows that we are still in the numerical conditions of the claim, and $i' < i$, so we can apply the induction hypothesis, except in the case that $d = m + 2$. In this case, either we are in one of the exceptions above, or $i = m + 1$, $j = 3$, $\mathbf{m}' = (m+2, 1^{m+3})$ and the claim follows from lemma 4.3.

m odd Write $m = 2\,t - 1$. By semicontinuity it is enough to see that there are no plane curves of degree d containing $Z_{K,\mathbf{m}}$ with K general in U_{t+1}. But for $K \in U_{t+1}$, unloading gives $Z_{K,\mathbf{m}} = Z_{K,(m+1,2^{i-t},1^{j+t})}$, so the result follows by the induction hypothesis again.

□

Remark that theorem 4.8 applies in particular when we have only double points, proper or infinitely near, and they are neither 5 nor less than 4. For the remaining cases the behaviour is also known (cf. [17]). Namely, one double point

has (obviously) maximal rank and the scheme of three double points (proper or infinitely near, in general position) has maximal rank; systems which fail to have maximal rank appear in degree 2 when there are two points and in degree 4 when there are five. In both cases there is an "unexpected" curve of the form $C = 2D$, where D is the straight line or the conic through the points, respectively. The cases of one point of multiplicity $m = 3, 4$ or 5 and i double points (again, proper or infinitely near) are also covered, except for $i < 2m/3+1$ and for the case $m = 4, i = 6$. All excepted cases involve less than eight points and were therefore also solved by B. Harbourne. The systems which do not have maximal rank are: $(3, 2)$ in degree 3, $(4, 2)$, $(4, 2^2)$ in degree 4, $(5, 2)$, $(5, 2^2)$ in degree 5, and $(4, 2^6)$ in degree 6.

We will now apply this result to find irreducible curves of low degree with tacnodes and cusps. We will use a form of Bertini's theorems slightly different from the usual ones. Given a linear system $\mathcal{L}$ of curves on S with no fixed part, the base points of $\mathcal{L}$ form a (usually non unibranched) cluster $\mathrm{BP}(\mathcal{L})$.

Proposition 4.9. *Let $\mathcal{L}$ be a linear system of curves on S with no fixed part. Then*

1. *General curves in $\mathcal{L}$ go sharply through $\mathrm{BP}(\mathcal{L})$.*
2. *If $\mathcal{L}$ is reducible then it is composed of the curves of a pencil.*

Proof. 2 and the fact that all singularities of general curves sit at the proper base points of $\mathcal{L}$ are standard. For a proof of 1, cf. [5, 7.2]. Cf. also Zariski's remark on the theorems of Bertini in [36]. □

Lemma 4.10. *Let $Z \subset Z'$ be two zero-dimensional subschemes of $\mathbb{P}^2$ such that $\operatorname{length} Z' = \operatorname{length} Z + 1$, and let d be a positive integer such that going through Z imposes independent conditions to curves of degree d. Then Z' imposes independent conditions to curves of degree $d + 1$.*

Proof. This is an easy application of the residual exact sequence of the Horace method. There is a unique point $p \in \mathbb{P}^2$ where the length of the component of Z' supported at p is bigger than that of Z. Let L be a general straight line through p, and consider the residual exact sequence

$$0 \longrightarrow \mathcal{I}_{Z''}(d) \longrightarrow \mathcal{I}_{Z'}(d+1) \longrightarrow \mathcal{I}_{Z'\cap L/L}(d+1) \longrightarrow 0$$

We have to prove that

$$H^1(\mathcal{I}_{Z'\cap L/L}(d+1)) = H^1(\mathcal{I}_{Z''}(d)) = 0\ .$$

As going through Z imposes independent conditions to curves of degree d and L is general, $\operatorname{length} Z \cap L \leq d + 1$, therefore $\operatorname{length} Z' \cap L \leq d + 2$ and $H^1(\mathcal{I}_{Z'\cap L/L}(d + 1)) = 0$. As by hypothesis $H^1(\mathcal{I}_Z(d)) = 0$, it will be enough to see that Z contains the residual scheme Z''. Let $I, I', I'' \subset \mathcal{O}_p$ be the ideals locally defining Z, Z' and Z'', and let $f \in \mathcal{O}_p$ be a local equation of L. $\operatorname{length} Z' = \operatorname{length} Z + 1$ implies that $I = I' + (g)$ for some $g \in \mathcal{O}_p$ with $g\mathfrak{m}_p \subset I'$, therefore $fI = fI' + (fg) \subset I'$ and $I \subset I'' = (I' : f)$, as wanted. □



Figure 3: Enriques diagrams of a tacnode cluster and an extended cusp cluster.

Corollary 4.11. *If $(K, \mathbf{m})$ is a weighted (not necessarily unibranched) cluster such that $Z_{K,\mathbf{m}}$ imposes independent conditions to curves of degree d, then general curves in $\mathcal{L} = \mathcal{L}_{d+1}(K, \mathbf{m})$ go sharply through $(K, \mathbf{m})$. Furthermore, if $(K, \mathbf{m})$ is not a single point of multiplicity $d+1$ then a general curve in $\mathcal{L}$ is irreducible.*

Proof. Because of lemma 4.10, $\mathcal{L}$ has no fixed part and $\mathrm{BP}(\mathcal{L}) = (K, \mathbf{m})$. By proposition 4.9 then, we only have to see that if $\mathcal{L}$ is composed of the curves of a pencil then $\mathrm{BP}(\mathcal{L})$ is a point with multiplicity $d+1$. Let $\mathcal{L}$ be composed of r curves of degree k in a pencil. Then $rk = d+1$ and

$$r = \ell_{d+1}(K, \mathbf{m}) = d + 2 + \ell_d(K, \mathbf{m}) \geq d+1$$

because $Z_{K,\mathbf{m}}$ imposes independent conditions to curves of degree d. Therefore $r = d+1$ and $k = 1$. A pencil of lines has a unique base point p so curves in $\mathcal{L}$ are composed of $d+1$ lines through p and $\mathrm{BP}(\mathcal{L})$ is as claimed. □

The cluster of infinitely near singular points of a tacnode of order t is a weighted unibranched cluster K of t free points with multiplicities $\mathbf{m} = (2^t)$. The scheme $Z_{K,\mathbf{m}}$ is called a tacnode scheme. The cluster of infinitely near singular points of a cusp of order n is a weighted unibranched cluster K of $n+2$ points, the last of which being satellite and the others free, with multiplicities $\mathbf{m} = (2^n, 1^2)$. We define an "extended" unibranched cluster $(K', \mathbf{m}')$ which has an additional free point taken with multiplicity one and call $Z_{K',\mathbf{m}'}$ a cusp scheme of order n. When this last point $p_{n+3} \in K'$ varies in $E_{n+2} \cong \mathbb{P}^1$ we obtain a flat family of cluster schemes, because of lemma 3.6. In the special position of p_{n+3} which makes it proximate to p_{n+1}, unloading gives $\delta(\mathbf{m}') =$

$(2^{n+1}, 0, 0)$ so any cusp scheme of order n can be specialized to the tacnode scheme of order $n+1$ given by the free points $p_1, \ldots, p_{n+1}$.

Theorem 4.12. *If*

$$\frac{d(d+1)}{6} \geq \sum_{i=1}^{\tau} t_i + \sum_{i=1}^{\nu} (n_i + 1) \geq 6$$

then there exists a reduced irreducible curve of degree d with τ tacnodes of orders $t_1, \ldots, t_\tau$ and ν cusps of orders $n_1, \ldots, n_\nu$ as its only singularities.

Proof. Consider a scheme

$$Z = T_1 \cup T_2 \cup \cdots \cup T_\tau \cup N_1 \cup N_2 \cup \cdots \cup N_\nu$$

where T_i is a tacnode scheme of order t_i supported at $p_i \in \mathbb{P}^2$ and N_i is a cusp scheme of order n_i supported at q_i, all of them having their points in general position. We claim that the linear system $\mathcal{L}_d(Z)$ of curves of degree d containing Z is nonempty and that a generic curve in it has a tacnode of order t_i at p_i, a cusp of order n_i at q_i and no other singularities.

Specializing the cusp schemes N_i to tacnode schemes $\bar{T}_i$ of order $n_i + 1$ we obtain a scheme $\bar{Z}$ in the conditions of theorem 4.8, therefore of maximal rank and length $3\left(\sum t_i + \sum (n_i + 1)\right)$. So by semicontinuity Z is also of maximal rank. The bound on d assures that Z imposes independent conditions to curves of degree $d-1$. So by 4.11 a general curve in $\mathcal{L}_d(Z)$ is irreducible and has no other singularities but the ones in p_i, q_i, which are tacnodes and cusps of the desired orders. □

Remark that in the hypotheses of theorem 4.12 we ask that $\sum t_i + \sum (n_i + 1) \geq 6$ in order to apply 4.8. In fact, the remarks we made after the proof of theorem 4.8 prove that the result holds also for $\sum t_i + \sum (n_i + 1) = 3, 4$. In case $\sum t_i + \sum (n_i + 1) = 5$, an ad–hoc reasoning can be used to prove the existence; however, for these small numbers of singularities the result is neither new nor significant, so we omit this.

We would like to point out that there are examples in [27] of curves F_k with one tacnode or cusp, of degree lower than the one given by 4.12. Namely, they have degree $d = 2k + 1$ and

$$\begin{cases} \text{a tacnode of order } \frac{2k^2+3k-1}{2} \text{ if } 2k^2 + 3k - 1 \text{ is even} \\ \text{a cusp of order } \frac{2k^2+3k-2}{2} \text{ if } 2k^2 + 3k - 1 \text{ is odd} \end{cases}$$

The author was informed by C. Lossen that he conjectures, after testing many particular cases, that these curves are irreducible and have no other singularity. He conjectures also that they do *not* satisfy the *T–smoothness* property (cf. [35]) which states that the variety of curves of degree d with that singularity is smooth of the expected dimension at F. Curves with tacnodes and cusps whose points are in general position (as those given by 4.12) do satisfy the T–smoothness property.

The reader may notice that after 4.11, any h^1–vanishing result for a class of cluster schemes can be exploited to obtain curves of low degree with the equisingularity type fixed by the clusters. In fact, the bounds obtained using 4.11 are

(slightly) sharper than those obtained using the somewhat more complicated reasoning of [14]. In particular, it is not difficult to extend theorem 4.12 to curves with tacnodes, cusps, and one different singularity.

Proposition 4.13. *Let S be an equisingularity class whose cluster of infinitely near singular points is unibranched and consists of one point of multiplicity m followed by k free double points. Define $M = k + \sum_{i=1}^{\tau} t_i + \sum_{i=1}^{\nu}(n_i+1) + \max\left(0, \frac{m^2-4m-6}{3}\right)$. If*

$$\frac{d(d+1)}{6} - \frac{m(m+1)}{6} \geq M \geq \frac{2}{3}m + 1$$

then there exists a reduced irreducible curve of degree d with one singularity of type S, τ tacnodes of orders $t_1, \ldots, t_\tau$ and ν cusps of orders $n_1, \ldots, n_\nu$ as its only singularities.

Proof. Analogous to the proof of 4.12. □

Remark also that in the proof of theorem 4.12 we proved that the union of general tacnode and cusp schemes has maximal rank, which is a case not included in theorem 4.8. It is easy to see that many other cases may be equally treated using the same techniques. For example,

Proposition 4.14. *The scheme of the cluster of infinitely near singular points of a D_k–singularity whose points are in general position has maximal rank, except in the two cases $k = 6, 7$.*

Proof. The cluster of infinitely near singular points of a D_k–singularity, with k even, is a weighted unibranched cluster with $k/2-1$ free points with multiplicities $\mathbf{m} = (3, 2^{k/2-2})$, so we are in a particular case of theorem 4.8.

If k is odd, then the cluster of infinitely near singular points of D_k is a weighted unibranched cluster with $r = (k+1)/2$ points, the last of which being satellite and the others free (that is, $K \in U_{r,r-1}$), with multiplicities $\mathbf{m} = (3, 2^{r-3}, 1^2)$. For simplicity, we assume $k \geq 13$ (or $r \geq 7$), as the cases with k small require special care. Let $U \subset P_{r,r-1} \cap P_3$ be the open subset where only p_3 and p_r are satellites (we specialize the third point to be proximate to the first). For $K \in U$, unloading gives $Z_{K,\mathbf{m}} = Z_{K,\mathbf{m}'}$ with $\mathbf{m}' = (4, 2^{r-4}, 1, 0, 0)$, so the length of $Z_{K,\mathbf{m}}$ is the same for K in U or in $U_{r,r-1}$, therefore $U \subset (P_{r,r+1})_{\mathbf{m}}$. As $U_{r,r-1}$ is dense in $U \subset (P_{r,r+1})_{\mathbf{m}}$, by semicontinuity it is enough to see that $Z_{K,\mathbf{m}} = Z_{K,\mathbf{m}'}$ has maximal rank for K general in U and the result follows from theorem 4.8, because the multiplicity of the last point is now 0 so we can assume K general in U_3. □

References

[1] Alexander, J. Hirschowitz, A., Interpolation on Jets *J. Alg.*, **192** (1997), 412-417.

[2] Barkats, D., Variétés des courbes planes à noeuds et à cusps, in: Peter E. Newstead, ed., *Algebraic Geometry,* Lect. Notes Pure App. Math. **200**, M. Dekker 1988, pp. 25-36.

[3] Brianon, J., Description de Hilb^n $\mathbb{C}\{x, y\}$, *Inventiones Math.* **41** (1977), 45-89.

[4] Casas-Alvero, E., Infinitely near imposed singularities and singularities of polar curves, *Math. Ann.* **287** (1990), 429-454.

[5] Casas-Alvero, E., *Singularities of plane curves* preprint Universitat de Barcelona (1997).

[6] Catalisano, M.V., Gimigliano, A. On curvilinear subschemes of $\mathbb{P}^2$ *J. Pure App. Algebra* **93** (1994), 1-14.

[7] Ciliberto, C. Miranda, R., Interpolation on Curvilinear Schemes, *J. Alg.* **203** (1998) 677-678.

[8] Ciliberto, C. Miranda, R., Degenerations of planar linear systems, *Journal Reine Ang. Math.* **501** (1998) 191-220.

[9] Ciliberto, C. Miranda, R., Linear systems of plane curves with base points of equal multiplicity, (1998) to appear in *Trans. A. M. S.*

[10] Enriques, F. - Chisini, O., *Lezioni sulla teoria geometrica delle equazioni e delle funzioni algebriche,* N. Zanichelli, Bologna 1915.

[11] Evain, L. *Collisions de trois gros points sur une surface algébrique*, thesis, Nice 1997.

[12] Evain, L., Une minoration du degr des courbes planes singularits imposes, *Preprint ENS Lyon,* **212** (1997), 1-17.

[13] Fogarty, J., Algebraic families on an algebraic surface, *Amer. J. Math* **10** (1968) 511-521.

[14] Greuel, G.M. - Lossen, C. - Shustin, E., Plane curves of minimal degree with prescribed singularities, *Inventiones Math.* **133** (1998), 539-580.

[15] Greuel, G.M. - Lossen, C. - Shustin, E., Castelnuovo function, zero-dimensional schemes and singular plane curves, *duke e–print* **9903179** (1999).

[16] Grothendieck, A. - Dieudonn, J., EGA IV, 4, *Elments de gomtrie algbrique,* Inst. Hautes Etudes Sci. Publ. Math. **32** (1967).

[17] Harbourne, B., Complete linear systems on rational surfaces, *Trans. A.M.S.* **289** (1985) 213-226.

[18] Harbourne, B., The geometry of rational surfaces and Hilbert functions of points in the plane, *Can. Math. Soc. Conf. Proc.* vol. **6** (1986) 95-111.

[19] Harbourne, B., Iterated blow-ups and moduli for rational surfaces, in: A. Holme and R. Speiser, eds., *Algebraic Geometry Sundance 1986*, LNM **1311**, Springer 1988, pp. 101-117.

[20] Harbourne, B., Rational surfaces with $K^2 > 0$, *Proc. A. M. S.* **124** (1996) 727-733.

[21] Harbourne, B., Anticanonical rational surfaces, *Trans. A.M.S.* **349** (1997) 1191-1208.

[22] Hartshorne, B., *Algebraic Geometry*, GTM 52, Springer 1977.

[23] Hirschowitz, A., La Mthode d'Horace pour l'interpolation a plusieurs variables, *Manuscripta Math.* **50** (1985), 337-388.

[24] Hirschowitz, A., Une conjecture pour la cohomologie des diviseurs sur les surfaces rationnelles generiques, *J. Reine Angew. Math* **397** (1989), 208-213.

[25] Iarrobino, A. Punctual Hilbert Schemes, *Mem. Amer. Math. Soc.* **188** (1977).

[26] Kleiman, S.L., Multiple-point formulas I: Iteration, *Acta Math.* **147** (1981), 13-49.

[27] Lossen, C., New asymptotics for the existence of plane curves with prescribed singularities, to appear in *Comm. in Algebra* (1999).

[28] Miranda, R., Linear systems of plane curves, *Notices of the A. M. S.* **46**, 2, (1999), 192-202.

[29] Nobile, A., Villamayor, O., Equisingular stratifications associated to families of planar ideals, *J. Alg.* **193** (1997), 239-259.

[30] Paxia, G., On flat families of fat points, *Proc. A. M. S.* **112** (1991), 19-23.

[31] Ran, Z., Curvilinear enumerative geometry, *Acta Math.* **155** (1985), 81-101.

[32] Ran, Z., Enumerative geometry of singular plane curves, *Inventiones Math.* **97** (1989), 447-469.

[33] Ran, Z., On the Nagata problem, *duke e–print* **9809101** (1998).

[34] Roé, J., On the existence of plane curves with imposed multiple points, *duke e–print* **9807066** (1998).

[35] Shustin, E., Smoothness of equisingular families of plane algebraic curves, *Int. Math. Res. Notices* (1997), 67-82.

[36] Zariski, O., *Algebraic Surfaces*, 2nd. suppl. ed., Ergebnisse 61, Springer 1971.

[37] Zariski, O., *Le problme des modules pour les branches planes*, Hermann 1986.